



DEFOLIATION IS AN IMPORTANT MEASURE

Ubaydullaev M.M.

Doctor of Philosophy in Agricultural Sciences, Head of the department, Fergana

Polytechnic Institute, Fergana, Uzbekistan

Sultonov S.T.

Researcher, Fergana Polytechnic Institute, Fergana, Uzbekistan

ABSTRACT: - The timing and depth of sowing should be determined by the soil conditions of each region and even each field. The most important is the timely cultivation of the rows.

KEYWORDS: Norm, term, soil, cultivator, a number of plants.

INTRODUCTION

The Republic of Uzbekistan is the most northern region of cotton-producing countries in the world. The climate here is continental, with rainy days lasting until March and even the first ten days of April. Therefore, the best time for sowing is during the second decade of April and the fifth day of May. Seeds should be sown correctly and in a timely manner with sufficient moisture to produce a high-quality cotton crop. When exposed to inadequate soil, most of the soil

decomposes and the rest grows poorly and sparingly [1-4].

THE MAIN PART

When determining the sowing time, it is important to take into account that the cotton is a warm plant, with a temperature of 20-25°C for normal growth and development [5-9]. Based on the results of many years of research and development experience of the research institutions, the sowing of seeds should be completed within the following timeframe:

“DEFOLIATION IS AN IMPORTANT MEASURE”

1. April 5-15 in Tashkent and Ferghana regions.
2. April 1-15 in Syrdarya, Jizzakh, Namangan, Andijan and Bukhara regions.
3. In mountainous areas of Andijan region and in Samarkand region April 5-20.
4. March 25 in the southern districts of Surkhandarya region Until April 10, April 1-15 in the Northern District.
5. In the southern regions of Kashkadarya from 25 March to 15 April, in the northern areas from 1 to 15 April.
6. It is recommended to plant in the Khorezm region and southern districts of the Karakalpak Autonomous Republic from April 10 to 25, and in the northern districts of the autonomous republic until April 15-30.

Seedless seeds are sown after 5-6 days in comparison with hairy seeds. Depending on the weather, these terms can be delayed for 2-3 days or backwards is possible. The timing and depth of sowing should be determined according to the soil conditions of each farm and even each field [10-17]. In our experience in grassy-soil soils of the Fergana region, the seeds of cotton seeds S8290 and S6775, according to the soil condition, were planted to a depth of 3-4 cm with a daily average soil temperature of 10 cm. When sowing seeds, it is necessary to take into account that the amount of residual fibre on the seeds should not exceed 0.8% in the medium-fibre types and 0.4% in the thin fibre varieties [18-21]. The seeds may not fall evenly from the sowing apparatus and the sprouts may be sparse. Providing fertilizers in most cotton fields is one of the most important tasks. According to the results of many years of research, from 7 to 10% of the annual fertilizer rate from cotton to weeding, nitrogen requires about 5-7% of phosphorus.

When cotton is sprouted, the soil will not be able to absorb the nitrogen and phosphorus it contains. That is why nitrogen, phosphorus and potassium fertilizers, given at the same time as sowing, have significant benefits for the rapid development of cotton. These fertilizers have a positive effect on the plant's faster absorption of mobile nitrogen, phosphorus and potassium in the soil. Based on the above scientific results, in our experience, we have 30–40 kg of urea per hectare and 40-50 kg of amorphous tractor cultivator per sow, with 6–8 rows per sowing row 10 cm deep and 10-12 cm deep [21-27]. In addition, the timely cultivation of cotton fields is one of the most important factors for achieving high yields. By increasing the inter-row cultivation of cotton, the activity of microorganisms in the soil will be improved, and the plant's micro-and macro-organisms will increase. As a result, the vitality of plant roots is improved through oxygen enrichment by air exchange in the soil. Rapid processing of young cotton plantations facilitates the development of young seedlings by providing porous soil and hot air. When working the rows, the sow is planted to a width of 60 cm, taking into account the age of the sprout.

If planted, appropriate soil mitigation and grazing facilities will be established.

CONCLUSION

During the second and subsequent processing, the working bodies may be installed 3-4, 5-6 cm deep, in both widths. It should be remembered that the number of cultivations is 6-7 times on light and sandy soils and 7-8 times on medium and heavy soils. 25-30 cm depth should be softened once. If this method is not applied (ie not deeply cultivated) before sowing, the soil will be interrupted by the reduction of moisture content in the soil before further cultivation. After 7-10 cm of soil, an artificial wall will be

“DEFOLIATION IS AN IMPORTANT MEASURE”

formed. This should not be allowed. From the experiments, it can be concluded that the fertilizer should be in proportion to the seeding rate and pay attention to the amount of residual fibres on the hairy seeds. Early cultivation allows the field to be well cleaned due to the loss of weeds during their early age.

REFERENCES

1. Esonzoda, S., Khalikova, Z., & Ibragimov, A. (2021). Determination of moisture and temperature of cotton from the drying drum with the IT. *International Engineering Journal For Research & Development*, 6(3), 7-7.
2. Odilzhanovich, T. K., Makhmudovna, N. M., & Odilzhanovich, I. A. (2021). The selection of the control parameter of the raw cotton electric sorter. *Innovative Technologica: Methodical Research Journal*, 2(11), 1-5.
3. NuraliQudratovich, S., AbdurahmonMuzaffarovich, E., & UlugbekTolibjonogli, T. (2020). To study the main factors influencing fiber quality in the process of sawdust separation and their interdependence. *European Journal of Molecular & Clinical Medicine*, 7(07), 2020.
4. Oripov, N., Komilov, J., Xolikova, Z., & Toshmirzaevk, O. Research on the Introduction of a Double-faced Improved Cotton Separator. *International Journal of Innovations in Engineering Research and Technology*, 7(12), 105-110.
5. Isaev, S. S., Yu, E., Oripov, N., & Xakimov, I. Study of the Effect on the Natural Characteristics of Fiber in the Process of Application of Cotton Processing Technology. *International Journal of Innovations in Engineering Research and Technology*, 7(12), 111-116.
6. Toshtemirov, Q. A., & Oripov, N. M. (2021). Improvement of ring spinning machine stretching equipment. *Innovative Technologica: Methodical Research Journal*, 2(10), 61-66.
7. Odilzhanovich, T. K., Odilzhanovich, I. A., & Makhmudovna, N. M. (2021). Analysis of FLUFF in the Process of Lintering of Seeds. *Central Asian journal of theoretical & applied sciences*, 2(11), 26-28.
8. Abdulhayevich, T. Q. (2021). Analysis of runners and spinners used in spinning machines. *Innovative Technologica: Methodical Research Journal*, 2(10), 34-37.
9. Shakhnoza, U., Mirpolat, K., Khasan, A., Rustam, A., Tulkin, O., & Islombek, N. (2021). Change of Quality Indicators of Fabric Fabrics. *Annals of the Romanian Society for Cell Biology*, 25(6), 2869-2874.
10. Nabiyev, Q. Q., Yaqubov, N. J., & Toshtemirov, K. A. (2020). Innovative technology in the production of clothing from natural fibers. *ACADEMICIA: An International Multidisciplinary Research Journal*, 10(11), 1186-1191.
11. Бекмирзаев, Ш., Саидмахамадов, Н., & Убайдуллаев, М. (2016). Получения Литье В Песчано-Глинистые Методом. *Теория и практика современной науки*, (6-1), 112-115.
12. Usmonov, J. M., Shakirov, S. M., Ubaydullayev, M. M., & Parmonov, S. O. (2021). Aluminum-based composition materials for processing aluminum scrap. *ACADEMICIA: An International Multidisciplinary Research Journal*, 11(8), 590-595.
13. Sharifjanovich, S. O. (2021, November). The Velocity Distribution over the Cross Section Pipes of Pneumatic

“DEFOLIATION IS AN IMPORTANT MEASURE”

- Transport Installations Cotton. In International Conference On Multidisciplinary Research And Innovative Technologies (Vol. 2, pp. 29-34).
14. Sharipjanovich, S. O., Umarali og, T. D., & Qizi, B. M. N. (2021). Current State And Analysis Of Equipment For Cleaning And Selection Of Seeds. *International Journal of Progressive Sciences and Technologies*, 29(2), 337-342.
 15. Ergashev, Y., Xusanova, S., & Axmadjonov, D. (2022). Analysis of the fiber quality of cotton varieties grown by region. *Gospodarka i Innowacje*, 21, 242-244.
 16. Ubaydullayev, M. M., Ne'matova, F. J., & Marufjonov, A. (2021). Determination of efficiency of defoliation in medium-fiber cotton varieties. *Galaxy International Interdisciplinary Research Journal*, 9(11), 95-98.
 17. Кодиров, З. З., Ирискулов, Ф. С., Пулатов, А., & Убайдуллаев, М. (2018). Electronic libraries as a fact of contemporary information landscape. *Экономика и социум*, (3), 629-633.
 18. Ubaydullaev, M. M. U., Askarov, K. K., & Mirzaikromov, M. A. U. Effectiveness of new defoliant. *Theoretical & applied science Учредители: Теоретическая и прикладная наука*, (12), 789-792.
 19. Eminov, S. O., & Хокимов, А. Е. (2021). Composite polymer materials for use in working bodies of cotton processing machines and mechanisms. *ISJ Theoretical & Applied Science*, 11 (103), 922-924.
 20. Zikirov, M. C., Qosimova, S. F., & Qosimov, L. M. (2021). Direction of modern design activities. *Asian Journal of Multidimensional Research (AJMR)*, 10(2), 11-18.
 21. Tashlanova, N. D. (2019). Development of critical thinking of students in universities. *Problems of modern science and education*, (11-2), 144.
 22. Каримов, Н. М., Абдусаттаров, Б. К., Махмудова, Г., & Саримсаков, О. Ш. (2021). Пневматическая транспортировка хлопка-сырца на хлопкозаводах. In *Инновационные Подходы В Современной Науке* (pp. 61-70).
 23. Сидиков, А. Х., Махмудова, Г., Каримов, А. И., & Саримсаков, О. Ш. (2021). Изучение движения частиц хлопка и тяжёлых примесей в рабочей камере пневматического очистителя. *Universum: технические науки*, (2-2 (83)).
 24. Odiljonovich, T. Q. (2021). About automation of loading and unloading of cotton raw materials at cotton factory stations. *ACADEMICIA: An International Multidisciplinary Research Journal*, 11(10), 2068-2071.
 25. Тешаев, Ф. Ж., & Убайдуллаев, М. М. (2020). Определение эффективных норм новых дефолиантов в условиях лугово-солончаковых почв Ферганской области при раскрытии коробочек 50-60% сортов хлопчатника с8290 и с6775. *Актуальные проблемы современной науки*, (5), 62-64.
 26. Ubaydullayev, Madaminjon Muminjonovich. (2021). *G'ozada defoliatsiya o'tkazishning maqbul me'yor va muddatlari. Monografiya. - Corresponding standards and terms of defliation of cotton. Monograph. - . Соответствующие нормы и сроки дефолиации хлопка. Монография. Zenodo.*

“DEFOLIATION IS AN IMPORTANT MEASURE”

<https://doi.org/10.5281/zenodo.57227>
21.

27. Mo'minovich, U. M. (2021). The Importance Of Planting And Processing Of Medium-Field Cotton Varieties Between Cotton Rows In Fergana Region. The American Journal of Agriculture and Biomedical Engineering, 3(09), 26-29.

“DEFOLIATION IS AN IMPORTANT MEASURE”