

**EUROPEAN INTERNATIONAL JOURNAL OF MULTIDISCIPLINARY
RESEARCH AND MANAGEMENT STUDIES****VOLUME04 ISSUE05**DOI: <https://doi.org/10.55640/eijmrms-04-05-25>

Pages: 149-156

**STUDY OF THE CONDITION OF PARODONT IN PERIODONTITIS IN FETAL WOMEN*****Abdulaxatov Abdugaffor Abdusattor Ugli****Clinic Resident Of The Department Of Orthopedic Dentistry, Uzbekistan****Islamova Nilufar Bustanovna****Phd, Assistant Department Of Orthopedic Dentistry Samarkand State Medical University Samarkand, Uzbekistan***ABOUT ARTICLE**

Key words: Gestation period, diseases of the hard tissue of the teeth and periodontal disease, periodontal disease, taste analyzer, chronic generalize periodontitis.

Received: 19.05.2024**Accepted:** 24.05.2024**Published:** 29.05.2024

Abstract: Pregnant women have the highest risk of dental diseases [3]. According to a number of authors, the prevalence of dental caries during the physiological period of pregnancy is 91.4%, in 90% of cases, diseases of the parodont tissue, mainly damage to healthy teeth with an acute course of the carious process, make up 38% of pregnant women. [5]. Secondary caries, development of the caries process, enamel hyperesthesia occur in 79% of pregnant women [11]. At the same time, the growth intensity of dental caries in terms of the absolute increase in the index of caries-fillers-removed teeth during pregnancy is 0.83 [2].

INTRODUCTION

In 50% of pregnant women and women in labor, gingivitis of pregnant women is observed in the normal period of pregnancy at 2-3 months of pregnancy. From the second half of pregnancy, the pathological process becomes more pronounced and often proceeds as general catarrhal or hypertrophic gingivitis and often develops pyogenic granuloma. With the development of pregnancy, periodont diseases develop continuously and only in the postpartum period does the clinical picture improve [4]. In the long run, gingivitis that occurs during pregnancy becomes chronic [9, 10].

Against the background of a change in reactivity in a pregnant woman and a decrease in the resistance of the body, foci of latent odontogenic infection can cause serious complications as a result of an exacerbation of the inflammatory process [7, 8].

The strongest strain of inflammatory events in the tissues of the Parodont occurs in the second trimester of pregnancy, and a sharp increase in the carisogenic condition in the oral cavity occurs in the third trimester [4, 2], which not only determines the optimal time for dental examinations. it is important to distinguish between the approach to programs for the prevention and treatment of the most important diseases of the oral cavity during pregnancy and in the postpartum period, as well as for this period of pregnancy.

A decisive role in the development of caries and periodontitis in pregnant women is played by hormonal substances produced by the placenta (somatomammotropin, progesterone, gonadotropin), changes in mineral and protein metabolism, immunological condition, oral microflora [1, 5].

It is not enough to talk about an increased risk of the appearance and development of periodontal diseases, as well as an increase in the intensity of dental caries in pregnant women; tires. This does not negate the importance of already identified factors that affect the appearance and development of dental caries, periodontal disease during pregnancy, but is aimed at competently combining diagnostic important known and newly identified criteria into a system of practical recommendations for identifying risk groups among pregnant women. dental health and their special monitoring.

Immunoprotective peptides of the biological environment, including oral fluid, are signs of local inflammatory intensity and are responsible for the implementation of innate immunity to microbes [6, 12]. In oral fluid, a whole complex of immunoprotective peptides is secreted, among which lactoferrin, catechidin LL-37 and a-defensin are secreted [14]. The introduction of immunoprotective peptides into algorithms and models for sorting the risk of developing dental caries during pregnancy extends the limits of existing recommendations in this direction.

Oral fluid and peripheral blood are used as biological agents in the study of systemic immunity in pregnant women with periodontal diseases [8]. However, despite the easy way to choose after the birth of a child and placenta, which is not associated with invasive manipulations, retroplacental blood and umbilical cord blood, which are fetal and maternal blood, are not used as biological agents. At the same time, the spectrum of inflammatory mediators, the determination of the mineral composition of retroplacental and umbilical blood allows to answer the questions of varying degrees of whether the inflammation of the periodontal tissues came with the "mother-placenta-fetus" system. Is it possible to

predict the appearance of caries of milk teeth in the future from the stage of birth? The study of these causal relationships enriches the existing scientific concept of the consequences of dental diseases for the body of the entire pregnant woman and develops new methods of prenatal prognosis of diseases.

Thus, in modern conditions, the organization of diagnosis, monitoring, prevention and prognosis of oral diseases of pregnant women and the provision of dental assistance to them remains an unresolved problem that determines the relevance of studies in this direction.

The purpose of this study is to optimize the system of dental care for pregnant women using clinical and laboratory indicators.

Materials and research methods. Dental examination of pregnant women with periodontal diseases and hard tissues for the study of clinical and immunological indicators in the development of periodontal diseases and hard tissues of the teeth, 25 people in the regional dental Polyclinic - the main group, as well as 15 people who were not during pregnancy. These patients were accepted as a comparison group.

Results and discussion. According to an analysis of research by some scientists, the incidence of dental caries in women during pregnancy is 90.8% and increases to 95% with preeclampsia [6]. In 39% of women, previously healthy teeth during pregnancy are damaged by acute caries lesions. According to many scientists, inflammatory and destructive lesions of periodontal tissues during pregnancy are observed in 65-90% of cases, and their prevalence in the prenatal period is 100% [10]. According to foreign authors, the incidence of damage to the periodontal tissue, starting with inflammation of the gums and ending with inflammation of the periodontal tissue, is 15-65%.

Lakoine M.O ishida. It has been found that during pregnancy, women have 35-98% gum inflammation, while periodontal inflammations are 7-22% [11]. Waget K. in the work of, a study with periodontitis in 345 pregnant women found that the depth of the pathological pocket lesion was mainly (38.7%) 4-6 mm [13]. The transition of dental caries to a complex form is due to its increase, re-developing caries occurs in 70% of pregnant women, while the increase in the caries process during pregnancy is 0.85% [5]. A characteristic sign of the development of caries in pregnant patients is the deep spread of the carious process to enamel and dentin, which is associated with damage to the highest percentage of teeth, including inflammation of the dental pulp [8].

80% enamel sensitivity in pregnant patients, hyperesthesia to mechanical, thermal and chemical factors of healthy teeth, the most common wedge-shaped defects, pathological wear of teeth of the vertical type [1].

Of the total number of pregnant women, 95% of pregnant women need dental treatment [7], orthopedics - 57%, emergency surgeries are performed by surgeons in 2.4% of patients [6].

In an analysis of 150 women during pregnancy, it was found that the indicators of increased tension of carious teeth increased by the end of the gestation period compared to the first trimester [7]. In particular, scientists found that during 8-12 weeks of gestation, the KPUz and KPU_p indices show approximately 10.9 ± 0.8 and 23.5 ± 0.9 , with larger values in 34-40 weeks. respectively 14.1 ± 0.9 and 26.1 ± 0.7 . The assessment of the hygienic index among women of the same category showed its poor picture and the appropriateness of the intermediate indicators of the gum index for the inflammatory process of moderate severity.

Zasloff M. and Tanida T. according to a combination of low levels of lysozyme with reduced concentrations of immunoglobulins, lactoferrin is associated with hypersensitivity to infectious pathogens. Thus, the amount of antimicrobial proteins in the oral fluid reflects the state of local non-specific protective factors of the oral mucosa.

Antimicrobial peptides are the first line of defense of Innate Immunity in all types of living beings, and the high importance of AMP is confirmed by their high content in circular neutrophils [1].

Studies have shown that amps are involved in an inflammatory response by producing chemoattractants for immune cells, including interleukin-8, and mobilizing immunocompetent T cells to attract neutrophils, as well as acting as enhancers for cell adhesion and later transepithelial cell migration. 12, 13, 14].

Recently, another structurally distinct subfamilies of γ -defensins have been identified in leukocytes of rhesus monkeys. In humans, γ -defensin secretion is blocked by mutations, but is conserved in primates [12].

REFERENCES

1. Alimjanovich R. J. et al. MINIMALLY INVASIVE METHODS OF TREATMENT OF UROLITHIASIS IN CHILDREN. REVIEW OF THE CURRENT STATUS OF THE ISSUES //JOURNAL OF BIOMEDICINE AND PRACTICE. – 2023. – T. 8. – №. 2.

2. Astanovich A. A. Comparative Analysis of the Stress-Strain State of the Lower Jaw with Different Splinting Systems in Localized Periodontitis of Middle Gravity by Finite Element Modeling. – 2023.
3. Astanovich, A. D. A., Alimdjanovich, R. J., Abdujamilevich, S. A., & Bakhriddnovich, T. A. (2021). The State of Periodontal Tissues in Athletes Engaged in Cyclic Sports. *Annals of the Romanian Society for Cell Biology*, 235-241.
4. Astanovich, A. A., & Ikrombekovna, A. N. (2024). IMPROVEMENT IN THE DIAGNOSIS OF CHRONIC PERIODONTITIS IN CHILDREN. *World Bulletin of Social Sciences*, 32, 68-70.
5. Asrorovna, X. N., Baxriddinovich, T. A., Bustanovna, I. N., Valijon O'g'li, D. S., & Qizi, T. K. F. (2021). Clinical Application Of Dental Photography By A Dentist. *The American Journal of Medical Sciences and Pharmaceutical Research*, 3(09), 10-13.
6. Bustanovna, I. N., & Sharipovna, N. N. (2023). Research cases in women after menopause clinical and morphological changes in oral organs and their analysis. *Journal of biomedicine and practice*, 8(3).
7. Bustanovna P. D. I. N. et al. Further Research the Features of the Use of Metal-Ceramic Structures in Anomalies of Development and Position of Teeth // *International Journal of Scientific Trends*. – 2024. – Т. 3. – №. 3. – С. 67-71.
8. Bustanovna, I. N. (2024). PATHOGENESIS OF PERIODONTAL DISEASE IN ELDERLY WOMEN. *Лучшие интеллектуальные исследования*, 21(3), 25-29.
9. Bustanovna, P. I. N. (2024). Research of the Structure of Somatic Pathology in Patients with Aphthous Stomatitis. *International Journal of Scientific Trends*, 3(3), 51-55.
10. Bustanovna, I. N. (2024). Complications Arising in the Oral Cavity after Polychemotherapy in Patients with Hemablastoses. *International Journal of Scientific Trends*, 3(3), 62-66.
11. Bustanovna, I. N. (2024). CLINICAL AND LABORATORY CHANGES IN PERIODONTITIS. *Journal of new century innovations*, 51(2), 58-65.
12. Bustanovna I. N. PATHOGENESIS OF PERIODONTAL DISEASE IN ELDERLY WOMEN // *Лучшие интеллектуальные исследования*. – 2024. – Т. 21. – №. 3. – С. 25-29.
13. Bustanovna, I. N. (2024). TO STUDY THE HYGIENIC ASSESSMENT OF THE CONDITION OF THE ORAL MUCOSA DURING ORTHOPEDIC TREATMENT. *Лучшие интеллектуальные исследования*, 21(1), 9-15.
14. Bustanovna, I. N., & Abdusattor o'g, A. A. A. (2024). Analysis of Errors and Complications in the Use of Endocal Structures Used in Dentistry. *International Journal of Scientific Trends*, 3(3), 82-86.

15. Bustanovna, I. N. (2024). The Effectiveness of the Use of the Drug " Proroot MTA" in the Therapeutic and Surgical Treatment of Periodontitis. *International Journal of Scientific Trends*, 3(3), 72-75.
16. Bustanovna, I. N. (2024). Hygienic Assessment of The Condition of The Oral Mucosa After Orthopedic Treatment. *International Journal of Scientific Trends*, 3(3), 56-61.
17. Bustanovna, I. N., & Abdusattor o'g, A. A. A. (2024). Analysis of Errors and Complications in the Use of Endocal Structures Used in Dentistry. *International Journal of Scientific Trends*, 3(3), 82-86.
18. Islamova, N. B. (2022). CHANGES IN PERIODONTAL TISSUES IN THE POSTMENOPAUSAL PERIOD. In *Стоматология-наука и практика, перспективы развития* (pp. 240-241).
19. Islamova, N. B., & Nazarova, N. S. (2023). IMPROVING THE DIAGNOSIS AND TREATMENT OF CHRONIC GENERALIZED PERIODONTITIS IN POSTMENOPAUSAL WOMEN. *Conferences*.
20. Islamova N. B., Sh N. N. STUDY OF CHANGES IN PERIODONTAL DISEASES IN POSTMENOPAUSAL WOMEN //Conferences. – 2023. – С. 15-17.
21. Исламова, Н. Б. (2024). Complications Arising in the Oral Cavity after Polychemotherapy in Patients with Hemablastosis. *International Journal of Scientific Trends*, 3(3), 76-81.
22. Исламова, Н. Б. (2024). ПАРОДОНТ КАСАЛЛИКЛАРИДА ОРГАНИЗМДАГИ УМУМИЙ ЎЗГАРИШЛАРНИ ТАҲЛИЛИ ВА ДАВОЛАШ САМАРАДОРЛИГИНИ ТАКОМИЛЛАШТИРИШ. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 43(7), 18-22.
23. Исламова, Н. Б., & Назарова, Н. Ш. (2023). СУРУНКАЛИ ТАРҚАЛГАН ПАРОДОНТИТ БИЛАН КАСАЛЛАНГАН ПОСТМЕНОПАУЗА ДАВРИДАГИ АЁЛЛАРНИНГ ПАРОДОНТ ТЎҚИМАСИНИНГ ДАВОЛАШ САМАРАДОРЛИГИ ОШИРИШ. *ЖУРНАЛ СТОМАТОЛОГИИ И КРАНИОФАЦИАЛЬНЫХ ИССЛЕДОВАНИЙ*, 4(2).
24. Исламова, Н. Б., & Исламов, Л. Б. (2021). Особенности развития и течения заболеваний полости рта при эндокринной патологии. *ББК*, 56, 76.
25. Исламова, Н., Хаджиметов, А., & Шакиров, Ш. (2015). Роль продуктов перекисного окисления липидов и противовоспалительных цитокинов крови в развитии заболеваний полости рта при гипотиреозе. *Журнал проблемы биологии и медицины*, (1 (82)), 41-44.
26. Иргашев, Ш. Х., & Исламова, Н. Б. (2021). Применение и эффективность энтеросгеля при лечении генерализованного пародонтита. In *Актуальные вопросы стоматологии* (pp. 305-310).

27. Исламова, Н. Б., Шамсиев, Р. А., Шомуродова, Х. Р., & Ахмедова, Ф. А. (2014). Состояние кристаллообразующей функции слюны при различных патологиях. In Молодежь и медицинская наука в XXI веке (pp. 470-471).
28. Исламов, Б. С., & Исламова, З. Б. (2020). БИОЛОГИЯ СЕМЯН КУЗИНИИ ТЕНЕВОЙ (COUSINIA UMBROSA BUNGE). In Современная наука: перспективы, достижения и инновации (pp. 39-47).
29. Исламова, Н. Б., & Чакконов, Ф. Х. (2021). Изменения в тканях и органах рта при эндокринных заболеваниях. In Актуальные вопросы стоматологии (pp. 320-326).
30. Исламова, Н. Б. (2016). Сравнительная оценка противовоспалительных цитокинов крови в развитии заболеваний полости рта при гипотиреозе. Наука в современном мире: теория и практика, (1), 41-44.
31. Иргашев, Ш., Норбутаев, А., & Исламова, Н. (2020). Эффективность энтеросгеля при лечении генерализованного пародонтита у ликвидаторов последствий аварии на чернойбыльской АЭС. Общество и инновации, 1(1/S), 656-663.
32. Исламова, Н. Б., & Назарова, Н. Ш. (2023, May). СОВЕРШЕНСТВОВАНИЕ ДИАГНОСТИКИ И ЛЕЧЕНИЯ ХРОНИЧЕСКОГО ГЕНЕРАЛИЗОВАННОГО ПАРОДОНТИТА У ЖЕНЩИН В ПЕРИОД ПОСТМЕНОПАУЗЫ. In Conferences (pp. 13-15).
33. Исламова, Н. Б., & Назарова, Н. Ш. (2023, May). СОВЕРШЕНСТВОВАНИЕ ДИАГНОСТИКИ И ЛЕЧЕНИЯ ХРОНИЧЕСКОГО ГЕНЕРАЛИЗОВАННОГО ПАРОДОНТИТА У ЖЕНЩИН В ПЕРИОД ПОСТМЕНОПАУЗЫ. In Conferences (pp. 13-15).
34. Исламова, Н. Б., & Назарова, Н. Ш. (2023). МЕТОДЫ ИССЛЕДОВАНИЯ ЗАБОЛЕВАНИЙ ПАРОДОНТА У ЖЕНЩИН, НАХОДЯЩИХСЯ В ПЕРИОДЕ ПОСТМЕНОПАУЗЫ. In АКТУАЛЬНЫЕ ВОПРОСЫ СТОМАТОЛОГИИ (pp. 334-338).
35. Maksudjanovna, M. G., & Bustanovna, I. Z. (2020, June). FIRST CELL OBSERVATIONS AND RESEARCH. In Archive of Conferences (Vol. 1, No. 1, pp. 142-143).
36. Назарова, Н., & Исламова, Н. (2022). Этиопатогенетические факторы развития заболеваний пародонта у женщин в периоде постменопаузы. Профилактическая медицина и здоровье, 1(1), 55-63.
37. Назарова, Н., & Исламова, Н. (2022). Assessment of clinical and morphological changes in the oral organs and tissues in post-menopause women. in Library, 22(1), 60-67.
38. Назарова, Н., & Исламова, Н. (2022). Postmenopauza davridagi ayollarda stomatologik kasalliklarining klinik va mikrobiologik ko'rsatmalari va mexanizmlari. in Library, 22(1), 204-211.

39. Ortikova N., Rizaev J. The Prevalence And Reasons Of Stomatophobia In Children //E-Conference Globe. – 2021. – С. 339-341.
40. Очилов, Х. У., & Исламова, Н. Б. (2024). Особенности артикуляции и окклюзии зубных рядов у пациентов с генерализованной формой повышенного стирания. SAMARALI TA'LIM VA BARQAROR INNOVATSIYALAR JURNALI, 2(4), 422-430.
41. Sharipovna, N. N., & Bustanovna, I. N. (2022). Assessment of clinical and morphological changes in the oral organs and tissues in post-menopause women. Frontline medical sciences and pharmaceutical journal, 2(05), 60-67.
42. Sulaymonova, Z. Z., & Islamova, N. B. (2023, May). TAKING IMPRESSIONS IN THE ORAL CAVITY AND THEIR REDUCTION. In Conferences (pp. 21-23).
43. Содикова, Ш. А., & Исламова, Н. Б. (2021). Оптимизация лечебно-профилактических мероприятий при заболеваниях пародонта беременных женщин с железодефицитной анемией. In Актуальные вопросы стоматологии (pp. 434-440).
44. Ризаев, Ж. А., & Ахмедов, А. А. (2023). GROWTH AND DEVELOPMENT OF GENERAL MEDICAL PRACTICE IN THE REPUBLIC OF UZBEKISTAN TO IMPROVE DENTAL CARE. ЖУРНАЛ СТОМАТОЛОГИИ И КРАНИОФАЦИАЛЬНЫХ ИССЛЕДОВАНИЙ, 4(3).
45. Xairullaevna O. N., Alimjanovich R. J. Improving the effectiveness of therapeutic and preventive measures by correcting psychoemotional stress in children at a dental appointment. – 2022.